



PRAKTIKUM PARASITOLOGI 1

Sodikin Kurniawan

www.sodiy.com.id

PERTEMUAN 6

H. nana, H. diminuta, T. saginata, T. solium, D. caninum dan D. Latum

Lecture prepared by Sodikin Kurniawan, Muhammadiyah University of Semarang




Cestode

Lecture prepared by Sodikin Kurniawan, Muhammadiyah University of Semarang






Classification of Cestodes – Man

Order	Adult worm seen in human intestine	Larval stage seen in humans
Pseudophyllidea	Diphyllobothrium latum, the fish tapeworm	<ul style="list-style-type: none"> • Spirometra mansoni
		<ul style="list-style-type: none"> • Spirometra theileri
		<ul style="list-style-type: none"> • Spirometra erinace
Cyclophyllidea	<ul style="list-style-type: none"> • Taenia saginata, the beef tapeworm • Taenia solium, the pork tapeworm • Hymenolepis nana, the dwarf tapeworm • Hymenolepis diminuta, the rat tapeworm (rare) • Dipylidium caninum, the double-pored dog tapeworm (rare) 	<ul style="list-style-type: none"> • Taenia solium, the pork tapeworm • Echinococcus granulosus, the dog tapeworm • Echinococcus multilocularis • Multiceps multiceps and other species

Tapeworm Morphology



Cacing Pita - *Tapeworm*

	<i>Taenia solium</i>	<i>Taenia saginata</i>	<i>Hymenolepis nana</i>	<i>Hymenolepis diminuta</i>	<i>Diphyllobothrium latum</i>
Heads					
	4 suckers 2 rows of hooks	4 suckers No hooks	4 suckers single row of 20–30 hooks	4 suckers No hooks	2 Suctorial grooves or bothria No suckers, No hooks
Proglottids					
	Longer than broad 7–12 uterine branches on each side	Longer than broad 15–30 uterine branches on each side	Broader than long	Broader than long	Broader than long Uterus coiled



Taenia sp.

Tenia saginata & Taenia solium

Lecture prepared by Sodikin Kurniawan, Muhammadiyah University of Semarang


Taenia sp.




Taxonomy

- *Cestoidea*
- *Order Cyclophyllidea* Lankester, 1877, *sedis mutabilis*
- *Family Taeniidae* Ludwig, 1886
- *Subfamily Taeniinae*^T Ludwig, 1886
- *Genus Taenia*^T Linnaeus, 1758
 - *Taenia solium* Linnaeus, 1758
 - [*Taenia saginata*] see *Taeniarhynchus saginata*

Taenia saginata - Beef tapeworm




Taenia solium - Pork tapeworm




Telur *Taenia sp.*

- Bentuk bulat
- Dinding tebal
- Garis radier pada dinding
- Berisi embrio hexacanth




Scolex & Proglottids




Uterus
15-30 vs 7-12

Genital pore




Suckers



Taenia saginata

Hooklets
(2 rows)

Suckers



Taenia solium

Scolex & Proglottids



-

15 - 30



Rostellum

Uterus



+

7 - 12



Hymenolepis sp.

Hymenolepis diminuta & Hymenolepis nana


Lecture prepared by Sodikin Kurniawan, Muhammadiyah University of Semarang

Hymenolepis sp.

Taxonomy

- *Cestoidea*
- *Order Cyclophyllidea* Lankester, 1877, *sedis mutabilis*
- *Family Hymenolepididae* Perrier, 1897
- *Genus Hymenolepis*^T Weinland, 1858
 - [*Hymenolepis nana*] see *Vampirolepis nana*
 - *Hymenolepis diminuta*

Hymenolepis nana - Dwarf tapeworm




H. Diminuta –

Rat tapeworm



- lebih besar dari *H. nana*
- Siklus mirip *H. Nana*
- Infeksi tidak sengaja
- Tidak menimbulkan gejala

Telur *Hymenolepis* sp.



- Bentuk bulat atau sedikit oval.
- Ukuran *H. nana* 70 – 85 μm X 60 – 80 μm , *H. diminuta* dengan kisaran ukuran 30–50 μm .
- *H. diminuta* 4-8 filamen polar di antara kedua membran

Telur *Hymenolepis* sp.





H. diminuta vs H. nana

Membrane


Polar filaments (4-8 filaments)

Kait (6-hooked)


Oncosphere / hexacanth embryo



H. nana vs *H. diminuta* – Scolex & Proglottid






H. nana vs *H. diminuta* – Scolex



Rostellum

H. nana vs *H. diminuta*





Diphyllobothrium latum



Lecture prepared by Sodikin Kurniawan, Muhammadiyah University of Semarang

Diphyllobothrium latum


Taxonomy

- *Cestoidea*
- *Order Diphyllobothriidea* Kuchta, Scholz, Brabec & Bray, 2008
- *Family Diphyllobothriidae*^T Lühe, 1910
- *Genus Diphyllobothrium*^T Cobbold, 1858
- *Diphyllobothrium latum* (Linnaeus, 1758)


Telur *D. latum*



Diphyllobothrium latum




D. latum – Scolex & Proglottid



Scolex (head) is spatulate or spoon-shaped



Genital pore



2 slit-like longitudinal sucking grooves (bothria)



Dipylidium Caninum



Lecture prepared by Sodikin Kurniawan, Muhammadiyah University of Semarang

Dipylidium caninum


Taxonomy

- *Cestoidea*
- *Order Cyclophyllidea* Lankester, 1877, *sedis mutabilis*
- *Family Dipylidiidae* Railliet, 1896
- *Genus Dipylidium*^T Leuckart, 1863
- *Dipylidium caninum* (Linnaeus, 1758)

Dipylidium Caninum




Mature proglottid





Egg



Dipylidium Caninum



Eggs packet containing *Dipylidium* eggs inside




D. caninum - Scolex



Rostellum

Dipylidium Caninum





TERIMA KASIH

Sodikin Kurniawan

www.sodiy.com.id

PERTEMUAN 6

H. nana, H. diminuta, T. saginata, T. solium, D. caninum dan D. Latum

Lecture prepared by Sodikin Kurniawan, Muhammadiyah University of Semarang